

# Does Chronic Smoking Influence Fibrinolytic Potential in Type 1 Diabetes Mellitus?

A.J. Simpson,<sup>\*1</sup> N.A. Booth<sup>2</sup>, N.R. Moore<sup>2</sup>, R.S. Gray<sup>1</sup>

<sup>1</sup>Department of Medicine, St. John's Hospital, Livingston EH54 6PP, Scotland, UK

<sup>2</sup>Department of Molecular and Cell Biology, University of Aberdeen, Aberdeen AB25 2ZD, Scotland, UK

Tissue-type plasminogen activator (t-PA) and plasminogen activator inhibitor (PAI-1) were studied in 18 smokers and 18 closely matched non-smokers, all of whom had Type 1 diabetes mellitus (DM). None of the patients had advanced complications of diabetes. The t-PA and PAI-1 antigen levels were measured in plasma before and after venous occlusion, and were normal in Type 1 diabetes regardless of smoking status. Platelet PAI-1 levels were also measured and were found to be normal both in smokers and non-smokers. In smokers with Type 1 DM, plasma PAI-1 was significantly correlated with triglycerides. The normal fibrinolytic potential found in smokers with diabetes contrasts starkly with the significantly elevated plasma PAI-1 reported in smokers without diabetes. In smokers, triglycerides may effect low levels of PAI-1 release into plasma; this process may be significantly augmented in the presence of smoking-induced insulin resistance. The lack of endogenous insulin release in Type 1 diabetes may obviate the characteristic rise in plasma PAI-1 found in smokers who do not have diabetes. © 1998 John Wiley & Sons, Ltd.

*Diabet. Med.* 15: 683–687 (1998)

**KEY WORDS** Type 1 diabetes; smoking; fibrinolysis; plasma; platelets; tissue-type plasminogen activator; plasminogen activator inhibitor type 1

Received 24 October 1997; revised 2 February 1998; accepted 15 March 1998

## Introduction

The possible contribution of impaired fibrinolysis to the pathogenesis of thrombotic arterial disease has received increasing attention in recent years. Elevated plasma levels of plasminogen activator inhibitor type 1 (PAI-1) have been described in a variety of conditions associated with thrombosis.<sup>1–3</sup>

Type 1 diabetes mellitus (DM) is associated with a significantly increased mortality attributable to cardiovascular disease.<sup>4</sup> A wide variety of haemostatic abnormalities have been described in the condition<sup>5,6</sup> and may predispose toward the increased risk of coronary artery disease. The fibrinolytic enzyme system has been extensively studied in Type 1 DM but results have been conflicting (reviewed by Gough and Grant<sup>7</sup>). However, difficulties in interpretation of the existing literature include differences in methodology and the fact that several studies did not control for the effects of cigarette smoking.

We recently demonstrated that plasma PAI-1 is significantly elevated in cigarette smokers without clinical evidence of arterial disease.<sup>8</sup> We were interested in

studying the effects of smoking on fibrinolysis in patients with uncomplicated diabetes in order to determine whether a similar pattern is observed to that in the population without diabetes, and whether abnormal fibrinolysis is detectable before the evolution of advanced complications in Type 1 DM.

We and others have previously demonstrated that most circulating PAI-1 (over 90 %) resides in platelets,<sup>9,10</sup> albeit largely in an inactive form. A number of qualitative and quantitative platelet abnormalities have been described in Type 1 DM.<sup>11,12</sup> We therefore aimed to examine the effects of smoking on the fibrinolytic potential of plasma and platelets from patients with Type 1 DM, without evidence of advanced microangiopathic complications.

## Patients and Methods

### Patients

Patients were identified from the computerized registry held at our diabetes clinic and were invited to participate in the study if they: were aged between 25 and 40;<sup>13,14</sup> had Type 1 DM; attended the clinic regularly; had no record of complications of diabetes at their last attendance (including normal fundoscopy through dilated pupils at least once during the previous year). Patients were

\*Correspondence to: Dr. J. Simpson, Rayne Laboratory, Medical School, University of Edinburgh, Teviot Place, Edinburgh EH8 9AG, UK. E-mail: a.j.simpson@ed.ac.uk

eligible for the smoking group if they had smoked at least 15 cigarettes per day for at least 5 years and for the non-smoking group if they had never smoked. Wherever possible, the smoking history was corroborated by a partner or friend.

All patients attended for study in the morning, with smokers being asked to abstain from smoking from midnight the night before.<sup>15</sup> Patients followed their usual morning routine in terms of morning insulin and breakfast; fibrinolysis is not significantly altered immediately following exogenous insulin<sup>16</sup> or meals.<sup>17</sup> A medical history was taken and physical examination performed. All patients supplied a mid-stream sample of urine (MSU) and a sample for dipstick urinalysis.

Exclusion criteria for the study included: symptoms of hypoglycaemia on the morning of attendance; history suggestive of claudication or sensory impairment; history suggestive of ischaemic heart disease or of deep venous thrombosis; pregnancy; use of aspirin, non-steroidal anti-inflammatory drugs or the oral contraceptive pill; hypertension (defined as mean systolic blood pressure  $\geq 160$  mmHg and or mean diastolic blood pressure  $\geq 90$  mmHg); clinical absence of any of the peripheral pulses; evidence of retinopathy greater than grade 1; clinical evidence of sensory neuropathy or abnormal reflexes; anything greater than trace protein or trace blood on dipstick urinalysis; evidence of urinary tract infection on MSU; serum creatinine above the local reference range; clinical evidence of intercurrent illness.

Thirty-six patients (18 smokers and 18 non-smokers) satisfied the entry criteria. Two patients had less than three microaneurysms per fundus at fundoscopy. No other patients had evidence of retinopathy. A trace of haematuria was noted on urinalysis in one non-smoker and three smokers. A trace of proteinuria was found in three non-smokers and two smokers. All patients gave informed, written consent. The study was approved by the local Ethics Committee.

## Protocol

The protocol employed has been described in detail previously.<sup>8</sup> In brief, blood samples were taken between 8 am and 10 am, in view of the known diurnal variation in fibrinolysis.<sup>18</sup> Samples for t-PA and PAI-1 estimation were collected onto an anticoagulant/antiplatelet mixture (Diatube H, Diagnostica Stago, Asnières-sur-Seine, France).<sup>19</sup> A standard venous occlusion test was then performed.<sup>8</sup> We have previously shown that the rise in platelet count accompanying venous occlusion closely reflects the haemoconcentration invariably associated with the test<sup>8</sup> and therefore a rise in platelet count was used to correct for haemoconcentration in this study. Standard techniques were used to prepare platelet rich plasma (PRP) and platelet free plasma (PFP).<sup>9</sup> PAI-1 in PFP represents the plasma pool of PAI-1, whilst subtracting PAI-1 in PFP from PAI-1 in PRP gives the platelet pool

of PAI-1. Dividing the platelet pool of PAI-1 by the platelet count in PRP gives the level of PAI-1 per platelet.

## Assays

T-PA antigen was assayed by enzyme-linked immunosorbent assay (ELISA) as described by MacGregor *et al.*<sup>20</sup> PAI-1 antigen was assayed by ELISA using purified PAI-1 as standard. The PAI-1 ELISA<sup>21</sup> was modified as described previously.<sup>9</sup> Biochemical and haematological parameters were measured using standard laboratory techniques.

## Statistics

When comparing variables from different groups, the Mann-Whitney U test was used. Confidence intervals for differences between medians were calculated as described by Campbell and Gardner.<sup>22</sup> Spearman's rank correlation test was used when variables were tested for correlation.

## Results

### Baseline Characteristics of Patients

The smoking and non-smoking groups were closely matched for all measured baseline variables (Table 1). The smoking group had an average estimated cumulative exposure to cigarettes of 18.5 pack-years (range 7–28.8).

### T-PA and PAI-1

No significant difference was found when comparing t-PA or PAI-1 antigen levels in non-smokers and smokers,

Table 1. Clinical, metabolic and anthropometric characteristics of smokers and non-smokers

	Smokers	Non-smokers
<i>n</i>	18	18
M:F ratio	14:4	15:3
Age (yr)	33 (25–39)	33.5 (26–39)
Duration of DM (yr)	4 (1.5–10)	6.5 (1.5–24.0)
Insulin (U day <sup>-1</sup> )	63 (34–94)	59 (25–108)
BMI (kg m <sup>-2</sup> )	24.9 (20.1–33.1)	26.1 (19.0–29.2)
Blood pressure		
Systolic (mm Hg)	125.5 (109–138)	122 (107–147)
Diastolic (mm Hg)	70 (60–86)	73 (50–87)
Glucose (mmol l <sup>-1</sup> )	15.2 (3.6–21.8)	13.2 (2.3–22.7)
HbA <sub>1c</sub> (%)	7.4 (5.1–9.7)	6.8 (3.8–9.6)
Triglycerides (mmol l <sup>-1</sup> )	1.16 (0.41–2.43)	0.81 (0.38–2.64)
Cholesterol (mmol l <sup>-1</sup> )	4.7 (3.3–7.2)	4.6 (3.4–6.8)
Fibrinogen (g l <sup>-1</sup> )	2.8 (1.9–4.0)	2.5 (1.7–4.2)
Albumin (g l <sup>-1</sup> )	40.5 (38–43)	41 (37–46)

Results expressed as median and range.

either before or after venous occlusion (Table 2). For t-PA pre-venous occlusion, the 95 % confidence interval for the difference in medians was from  $-1.0$  to  $+1.8$  ng ml $^{-1}$ , and for PAI-1 from  $-9.5$  to  $+3.6$  ng ml $^{-1}$ . The increase in t-PA after venous occlusion exceeded that due to haemoconcentration in both groups, but the increment in each was similar, with no statistically significant difference observed. No significant difference was found between the levels of platelet PAI-1 in the two groups.

### Correlation between t-PA, PAI-1 and baseline variables

Plasma t-PA and PAI-1 levels were closely correlated both among non-smokers ( $r = 0.61$ ,  $p < 0.05$ ) and smokers ( $r = 0.63$ ,  $p < 0.01$ ). Among the smoking group, serum triglycerides correlated significantly with plasma PAI-1 ( $r = 0.49$ ,  $p < 0.05$ ) and t-PA ( $r = 0.54$ ,  $p < 0.05$ ). Neither PAI-1 nor t-PA correlated significantly with triglycerides in the non-smoking group.

### Discussion

Our findings show that chronic smoking has little, if any, effect on the levels of t-PA or PAI-1 in patients with Type 1 DM of relatively short duration and without advanced microangiopathy. A previous study has described significantly elevated PAI activity in smokers with much longer histories of diabetes, many of whom had varying degrees of diabetic complications<sup>23</sup> which may have affected fibrinolysis.<sup>24–26</sup> Moreover, in that study,<sup>23</sup> as with several previous studies on fibrinolysis and diabetes, blood was collected onto citrate alone, raising the possibility that some of the measured PAI in plasma had leaked from platelets *in vitro*.<sup>19</sup>

The current study was designed to minimize the

potential for diabetic complications to confound any observed effect of smoking on fibrinolysis. The finding of essentially normal t-PA and PAI-1 antigen levels in Type 1 DM patients without advanced complications broadly supports the data generated by groups who have subdivided patients according to the degree of diabetic complications,<sup>25,27–29</sup> although one group has described significantly reduced t-PA and PAI-1 antigen levels in uncomplicated diabetes.<sup>30</sup>

One of the main differences when comparing our study with previous work concerns the duration of diabetes in our patient groups being generally shorter than in other studies. In this context it is reassuring that evidence exists to support the prolonged maintenance of normal fibrinolysis in the setting of uncomplicated disease.<sup>24,31</sup>

It seems safe to conclude that the impairment of fibrinolysis associated with smoking is circumvented in Type 1 DM; that levels of the principal activator and inhibitor of fibrinolysis are normal in uncomplicated Type 1 DM, and that breakfast and exogenous insulin do not significantly alter t-PA or PAI-1 levels.

The response to venous occlusion was essentially normal in our patients with diabetes, consistent with findings elsewhere.<sup>24,28,29,32</sup> Our findings suggest that smoking does not significantly compromise non-specific endothelial cell function in diabetes prior to the development of advanced complications.

Surprisingly little information has previously been generated regarding platelet PAI-1. We found that the platelet pool of PAI-1 was quantitatively normal in smokers and non-smokers with diabetes, echoing the resistance of the platelet pool to change in a variety of conditions.<sup>33</sup>

We previously described the effects of chronic smoking in a healthy population, and found that fibrinolytic potential in plasma was significantly impaired in smokers due to an elevation in PAI-1,<sup>8</sup> in support of other published work.<sup>34,35</sup> The protocol for the current study was virtually identical in all respects to that employed in our population without diabetes, with the exception that patients with diabetes were allowed their usual breakfast and insulin on the morning of study. The groups of smokers and non-smokers with diabetes described here, and the respective groups without diabetes, were extremely similar with regard to baseline characteristics.<sup>8</sup> The previously described healthy non-smokers (in whom fibrinolysis can be considered normal) and healthy smokers therefore represent closely matched healthy control groups for the non-smokers and smokers with diabetes, respectively. A summary of data from the subjects without diabetes<sup>8</sup> and patients with diabetes is shown in Table 3.

The lack of association between smoking and reduced fibrinolytic potential in patients with Type 1 DM, contrasting with the pattern in subjects without diabetes, suggests a critical role for insulin secretion in mediating increased PAI-1 in smokers without diabetes. Smoking

Table 2. T-PA and PAI-1 in plasma and platelets

	Non-smokers	Smokers
Plasma t-PA antigen (ng ml $^{-1}$ )		
Before VO	4.4 (1.5–8.3)	3.8 (1.5–8.3)
After VO	19.2 (6.9–64.2)	15.6 (5.6–59.4)
Plasma PAI-1 antigen (ng ml $^{-1}$ )		
Before VO	10.2 (2.5–87.4)	14.3 (2.8–52.0)
After VO	13.9 (4.6–153.6)	20.2 (6.5–71.8)
Plasma t-PA:PAI-1 ratio		
Before VO	0.44 (0.07–2.60)	0.36 (0.07–1.00)
After VO	1.32 (0.34–6.80)	0.97 (0.18–2.15)
PAI-1 per platelet [ng PAI-1 (10 $^6$ platelets) $^{-1}$ ]	0.34 (0.15–1.39)	0.31 (0.14–0.92)

Results expressed as median (range). VO, venous occlusion.

Table 3. T-PA and PAI-1 levels: comparison of levels in smokers and non-smokers with or without diabetes

	Non-smokers with diabetes	Smokers with diabetes	Non-smokers without diabetes	Smokers without diabetes
<i>n</i>	18	18	18	18
Plasma PAI-1 (ng ml <sup>-1</sup> )	10.2 <sup>b</sup> 2.5–87.4	14.3 <sup>b</sup> 2.8–52.0	14.2 <sup>a</sup> 3.6–83.1	34.6 3.8–182.5
Plasma t-PA (ng ml <sup>-1</sup> )	4.4 1.5–8.3	3.8 <sup>a</sup> 1.5–8.3	5.0 1.6–17.5	6.2 1.6–24.6
Plasma t-PA:PAI-1 ratio	0.44 <sup>c</sup> 0.07–2.60	0.36 <sup>a</sup> 0.07–1.00	0.30 0.08–1.62	0.16 0.04–0.67
Platelet PAI-1 [ng PAI-1 (10 <sup>6</sup> platelets) <sup>-1</sup> ]	0.34 0.15–1.39	0.31 0.14–0.92	0.38 0.13–0.67	0.39 0.10–0.92

Results are expressed as median and range. For any given parameter measured, the only statistically significant differences were found when comparing 'smokers without diabetes' with other groups.

<sup>a</sup>*p*<0.05 compared with 'smokers without diabetes'; <sup>b</sup>*p*<0.005 compared with 'smokers without diabetes'; <sup>c</sup>*p*<0.001 compared with 'smokers without diabetes'.

Data relating to subjects without diabetes has been reported previously,<sup>8</sup> as described in the Discussion.

is recognized to cause insulin resistance,<sup>36,37</sup> and thus hyperinsulinaemia, in subjects without diabetes. Against this background, smoking can stimulate triglyceride production,<sup>38</sup> and multivariate analysis has shown that plasma PAI-1 levels correlate with VLDL concentrations rather than insulin concentration or measures of insulin sensitivity.<sup>39,40</sup> In the population with diabetes, as was found in the population without diabetes,<sup>8</sup> triglycerides correlated with plasma PAI-1 in smokers, but not in non-smokers. However it seems unlikely that triglycerides are solely responsible for the elevation in PAI-1 in smokers without diabetes, as no significant difference was found between triglyceride levels in smokers with and without diabetes, while PAI-1 levels were significantly higher in subjects without diabetes (Table 3). From the available evidence it may be inferred that smoking, via triglycerides/VLDLs, induces a low level of PAI-1 release which, in subjects without diabetes, is significantly augmented (either directly, or indirectly via VLDLs or triglycerides) by insulin resistance/hyperinsulinaemia. This hypothesis contends that such augmentation cannot be effected in diabetic subjects, in whom endogenous insulin release is conspicuously absent.

In conclusion we consider that in the absence of advanced complications, circulating fibrinolytic potential is normal in Type 1 DM, and that this situation is not altered by chronic smoking. The radically different response to smoking in patients with diabetes from that in smokers without diabetes offers important potential insights into the regulation of PAI-1 release.

## Acknowledgements

We are grateful for the helpful advice and access to facilities provided during the compilation of this study by the following people at St John's Hospital, Livingston: I. Okell and M. Scott (Diabetes Clinic); M. Cook, Consultant Haematologist, and D. McCullough, Consultant Clinical Chemist. We are also indebted to G. Murray,

Professor of Medical Statistics, University of Edinburgh, for advice relating to statistical analysis.

## References

1. Sprengers ED, Kluft C. Plasminogen activator inhibitors. *Blood* 1987; **69**: 381–387.
2. Booth NA. The natural inhibitors of fibrinolysis. In: Bloom AL, Forbes CD, Thomas DP, Tuddenham EGD, eds. *Haemostasis and Thrombosis*, 3rd edn. Edinburgh: Churchill Livingstone, 1994: 699–717.
3. Juhan-Vague I, Alessi MC. Plasminogen activator inhibitor 1 and atherothrombosis. *Thrombosis and Haemostasis* 1993; **70**: 138–143.
4. Kannel WB, McGee DL. Diabetes and cardiovascular disease: the Framingham study. *J Am Med Assoc* 1979; **241**: 2035–2038.
5. Fuller JH, Keen H, Jarrett RJ, Omer T, Meade TW, Chakrabarti R, North WRS, Stirling Y. Haemostatic variables associated with diabetes and its complications. *Br Med J* 1979; **2**: 964–966.
6. Dubrey SW, Reaveley DR, Seed M, Lane DA, Ireland H, O'Donnell M, O'Connor B, Noble MI, Leslie DG. Risk factors for cardiovascular disease in IDDM. A study of identical twins. *Diabetes* 1994; **43**: 831–835.
7. Gough SCL, Grant PJ. The fibrinolytic system in diabetes mellitus. *Diabetic Med* 1991; **8**: 898–905.
8. Simpson AJ, Gray RS, Moore NR, Booth NA. The effects of chronic smoking on the fibrinolytic potential of plasma and platelets. *Br J Haematol* 1997; **97**: 208–213.
9. Booth NA, Simpson AJ, Croll AM, Bennett B, MacGregor IR. Plasminogen activator inhibitor (PAI-1) in plasma and platelets. *Br J Haematol* 1988; **70**: 327–333.
10. Declerck PJ, Alessi MC, Verstreken MV, Kruithof EKO, Juhan-Vague I, Collen D. Measurement of plasminogen activator inhibitor 1 in biologic fluids with a murine monoclonal antibody-based enzyme-linked immunosorbent assay. *Blood* 1988; **71**: 220–225.
11. Colwell JA, Winocour PD, Halushka PV. Do platelets have anything to do with diabetic microvascular disease? *Diabetes* 1983; **32** (suppl 2): 14–19.
12. O'Donnell MJ, Le Guen CA, Lawson N, Gyde OHB, Barnett AH. Platelet behaviour and haemostatic variables in Type 1 (insulin-dependent) diabetic patients with and without albuminuria. *Diabetic Med* 1991; **8**: 624–628.



13. Hashimoto Y, Kobayashi A, Yamazaki N, Sugawara Y, Takada Y, Takada A. Relationship between age and plasma t-PA, PA-inhibitor and PA activity. *Thrombosis Res* 1987; **46**: 625–633.
14. Gleeup G, Winther K. The effect of ageing on platelet function and fibrinolytic activity. *Angiology* 1995; **46**: 715–718.
15. Allen RA, Kluft C, Brommer EJP. Acute effect of smoking on fibrinolysis: increase in the activity level of circulating extrinsic (tissue-type) plasminogen activator. *Eur J Clin Invest* 1984; **14**: 354–361.
16. Potter van Loon BJ, de Bart ACW, Radder JK, Frölich M, Kluft C, Meinders AE. Acute exogenous hyperinsulinaemia does not result in elevation of plasma plasminogen activator inhibitor (PAI-1) in humans. *Fibrinolysis* 1990; **4** (suppl 2): 93–94.
17. Hendriks HFJ, Veenstra J, Velthuis-te Wierik EJM, Schaafsma G, Kluft C. Effects of moderate dose of alcohol with evening meal on fibrinolytic factors. *Br Med J* 1994; **308**: 1003–1006.
18. Kluft C, Verheijen JH, Rijken DC, Chang GTG, Jie AFH, Onkelinx C. Diurnal fluctuations in the activity of the fast-acting t-PA inhibitor. In: Davidson JF, Donati MB, Coccheri S, eds. *Progress in Fibrinolysis*, vol. VII. Edinburgh: Churchill Livingstone, 117–119.
19. Juhan-Vague I, Alessi MC, Fossat C, Declerck PJ, Kruithof EKO. Plasma determination of plasminogen activator inhibitor 1 antigen must be performed in blood collected on antiplatelet/anticoagulant mixture. *Thrombosis and Haemostasis* 1987; **58**: 1096.
20. MacGregor IR, MacDonald S, Dawes J, Micklem LR, James K. A monoclonal antibody enzyme linked immunosorbent assay directed towards a fibrin binding region of tissue-type plasminogen activator. *Fibrinolysis* 1987; **1**: 247–252.
21. MacGregor IR, Booth NA. An enzyme-linked immunosorbent assay (ELISA) used to study the cellular secretion of endothelial plasminogen activator inhibitor (PAI-1). *Thrombosis and Haemostasis* 1988; **59**: 68–72.
22. Campbell MJ, Gardner MJ. Calculating confidence intervals for some non-parametric analyses. In: Gardner MJ, Altman DG, eds. *Statistics with Confidence—Confidence Intervals and Statistical Guidelines*, 1st edn. London: British Medical Journal Publications, 1989: 71–79.
23. Nilsson TK, Lithner F. Glycaemic control, smoking habits and diabetes duration affect the extrinsic fibrinolytic system in type 1 diabetic patients but microangiopathy does not. *Acta Med Scand* 1988; **224**: 123–129.
24. Haitas B, Barnes AJ, Cederholm-Williams SA, Moore J, Shogry MEC, Turner RC. Abnormal endothelial release of fibrinolytic activity and fibronectin in diabetic microangiopathy. *Diabetologia* 1984; **27**: 493–496.
25. Jensen T, Feldt-Rasmussen B, Bjerre-Knudsen J, Deckert T. Features of endothelial dysfunction in early diabetic nephropathy. *Lancet* 1989; **1**: 461–463.
26. Lee P, Jenkins A, Bourke C, Santamaria J, Paton C, Janus E, Best J. Prothrombotic and antithrombotic factors are elevated in patients with type 1 diabetes complicated by microalbuminuria. *Diabetic Med* 1993; **10**: 122–128.
27. Mahmoud R, Raccah D, Alessi MC, Aillaud I, Juhan-Vague I, Vague P. Fibrinolysis in insulin dependent diabetic patients with or without nephropathy. *Fibrinolysis* 1992; **6**: 105–109.
28. Vicari AM, Viganó D'Angelo S, Testa S, Comi G, Galardi G, Orsi E, D'Angelo A. Normal tissue plasminogen activator and plasminogen activator inhibitor activity in plasma from patients with type 1 diabetes mellitus. *Horm Metab Res* 1992; **24**: 516–519.
29. Walmsley D, Hampton KK, Grant PJ. Contrasting fibrinolytic responses in type 1 (insulin-dependent) and type 2 (non-insulin-dependent) diabetes. *Diabetic Med* 1991; **8**: 954–959.
30. Ostermann H, Tschöpe D, Greber W, Meyer-Rüsenberg H-M, van de Loo, J. Enhancement of spontaneous fibrinolytic activity in diabetic retinopathy. *Thrombosis and Haemostasis* 1992; **68**: 400–403.
31. McLaren M, Jennings PE, Forbes CD, Belch JFF. Fibrinolytic response to venous occlusion in diabetics with and without microangiopathy compared to normal age and sex matched controls. *Fibrinolysis* 1990; **4** (suppl 2): 95–96.
32. Wieczorek I, Pell ACH, McIver B, MacGregor IR, Ludlam CA, Frier BM. Coagulation and fibrinolytic systems in type 1 diabetes: effects of venous occlusion and insulin-induced hypoglycaemia. *Clinical Science* 1993; **84**: 79–86.
33. Simpson AJ, Booth NA, Moore NR, Bennett B. The platelet and plasma pools of plasminogen activator inhibitor (PAI-1) vary independently in disease. *Br J Haematol* 1990; **75**: 543–548.
34. Belch JFF, McArdle BM, Burns P, Lowe GDO, Forbes CD. The effects of acute smoking on platelet behaviour, fibrinolysis and haemorrhage in habitual smokers. *Thrombosis and Haemostasis* 1984; **51**: 6–8.
35. Kjaeldgaard A, Larsson B. Long-term treatment with combined oral contraceptives and cigarette smoking associated with impaired activity of tissue plasminogen activator. *Acta Obstet Gynecol Scand* 1986; **65**: 219–222.
36. Facchini FS, Hollenbeck CB, Jeppesen J, Chen Y-DI, Reaven GM. Insulin resistance and cigarette smoking. *Lancet* 1992; **339**: 1128–1130.
37. Potter van Loon BJ, Kluft C, Radder JK, Blankenstein, MA, Meinders AE. The cardiovascular risk factor plasminogen activator inhibitor type 1 is related to insulin resistance. *Metabolism: Clinical and Experimental* 1993; **42**: 945–949.
38. Craig WY, Palomaki GE, Haddow JE. Cigarette smoking and serum lipid and lipoprotein concentrations; an analysis of published data. *Br Med J* 1989; **298**: 784–788.
39. Asplund-Carlson A, Hamsten A, Wiman B, Carlson LA. Relationship between plasma plasminogen activator inhibitor-1 activity and VLDL triglyceride concentration, insulin levels and insulin sensitivity: studies in randomly selected normo- and hypertriglyceridaemic men. *Diabetologia* 1993; **36**: 817–825.
40. Mykkanen L, Ronnema T, Marniemi J, Haffner SM, Bergman R, Laakso M. Insulin sensitivity is not an independent determinant of plasma plasminogen activator inhibitor-1 activity. *Arteriosclerosis and Thrombosis* 1994; **14**: 1264–1271.